

**[www.batdangthuc.net](http://www.batdangthuc.net)**

Happy New Year 2008

Chuc Mung Nam Moi 2008

*Vietnam Inequality Forum - VIF -*

*www.batdangthuc.net*

★★★★★

**Ebook Written by:**

**VIF Community**

*User Group: All*

★★★

*This product is created for educational purpose.  
Please don't use it for any commercial purpose  
unless you got the right of the author. Please  
contact [www.batdangthuc.net](http://www.batdangthuc.net) for more details.*

*Dien Dan Bat Dang Thuc Viet Nam*

*www.batdangthuc.net*

★★★★★

**Editors**

*Dien Dan Bat Dang Thuc Viet Nam*

★★★

*Bài Viet Nay (cung voi file PDF di kem) duoc tao ra vi muc dich giao duc. Khong duoc su dung ban EBOOK nay duoi bat ky muc dich thuong mai nao, tru khi duoc su dong y cua tac gia. Moi chi tiet xin lien he: [www.batdangthuc.net](http://www.batdangthuc.net).*

# *Dien Dan Bat Dang Thuc Viet Nam*

[www.batdangthuc.net](http://www.batdangthuc.net)

★★★★★

## **Contributors Of The Book**

- ★ **Editor. Pham Kim Hung (hungkhtn)**  
Admin, VIF Forum, Student, Stanford University
- ★ **Editor. Nguyen Manh Dung (NguyenDungTN)**  
Super Mod, VIF Forum, Student, Hanoi National University
- ★ **Editor. Vu Thanh Van (VanDHHK)**  
Moderator, VIF Forum, Student, Hue National School
- ★ **Editor. Duong Duc Lam (dduclam)**  
Super Moderator, VIF Forum, Student, Civil Engineering University
- ★ **Editor. Le Thuc Trinh (pi3.14)**  
Moderator, VIF Forum, Student, High School
- ★ **Editor. Nguyen Thuc Vu Hoang (zaizai)**  
Super Moderator, VIF Forum, Student, High School
- ★ **Editors. And Other VIF members who help us a lot to complete this version**

# Inequalities From 2007 Mathematical Competition Over The World



**Example 1 (Iran National Mathematical Olympiad 2007).** Assume that  $a, b, c$  are three different positive real numbers. Prove that

$$\left| \frac{a+b}{a-b} + \frac{b+c}{b-c} + \frac{c+a}{c-a} \right| > 1.$$

**Example 2 (Iran National Mathematical Olympiad 2007).** Find the largest real  $T$  such that for each non-negative real numbers  $a, b, c, d, e$  such that  $a + b = c + d + e$ , then

$$\sqrt{a^2 + b^2 + c^2 + d^2 + e^2} \geq T(\sqrt{a} + \sqrt{b} + \sqrt{c} + \sqrt{d} + \sqrt{e})^2.$$

**Example 3 (Middle European Mathematical Olympiad 2007).** Let  $a, b, c, d$  be positive real numbers with  $a + b + c + d = 4$ . Prove that

$$a^2bc + b^2cd + c^2da + d^2ab \leq 4.$$

**Example 4 (Middle European Mathematical Olympiad 2007).** Let  $a, b, c, d$  be real numbers which satisfy  $\frac{1}{2} \leq a, b, c, d \leq 2$  and  $abcd = 1$ . Find the maximum value of

$$\left(a + \frac{1}{b}\right) \left(b + \frac{1}{c}\right) \left(c + \frac{1}{d}\right) \left(d + \frac{1}{a}\right).$$

**Example 5 (China Northern Mathematical Olympiad 2007).** Let  $a, b, c$  be side lengths of a triangle and  $a + b + c = 3$ . Find the minimum of

$$a^2 + b^2 + c^2 + \frac{4abc}{3}.$$

**Example 6 (China Northern Mathematical Olympiad 2007).** Let  $\alpha, \beta$  be acute angles. Find the maximum value of

$$\frac{(1 - \sqrt{\tan \alpha \tan \beta})^2}{\cot \alpha + \cot \beta}.$$

**Example 7 (China Northern Mathematical Olympiad 2007).** Let  $a, b, c$  be positive real numbers such that  $abc = 1$ . Prove that

$$\frac{a^k}{a+b} + \frac{b^k}{b+c} + \frac{c^k}{c+a} \geq \frac{3}{2},$$

for any positive integer  $k \geq 2$ .

**Example 8 (Croatia Team Selection Test 2007).** Let  $a, b, c > 0$  such that  $a + b + c = 1$ . Prove that

$$\frac{a^2}{b} + \frac{b^2}{c} + \frac{c^2}{a} \geq 3(a^2 + b^2 + c^2).$$

**Example 9 (Romania Junior Balkan Team Selection Tests 2007).** Let  $a, b, c$  three positive reals such that

$$\frac{1}{a+b+1} + \frac{1}{b+c+1} + \frac{1}{c+a+1} \geq 1.$$

Show that

$$a + b + c \geq ab + bc + ca.$$

**Example 10 (Romania Junior Balkan Team Selection Tests 2007).** Let  $x, y, z \geq 0$  be real numbers. Prove that

$$\frac{x^3 + y^3 + z^3}{3} \geq xyz + \frac{3}{4}|(x-y)(y-z)(z-x)|.$$

**Example 11 (Yugoslavia National Olympiad 2007).** Let  $k$  be a given natural number. Prove that for any positive numbers  $x, y, z$  with the sum 1 the following inequality holds

$$\frac{x^{k+2}}{x^{k+1} + y^k + z^k} + \frac{y^{k+2}}{y^{k+1} + z^k + x^k} + \frac{z^{k+2}}{z^{k+1} + x^k + y^k} \geq \frac{1}{7}.$$

**Example 12 (Cezar Lupu & Tudorel Lupu, Romania TST 2007).** For  $n \in \mathbb{N}, n \geq 2, a_i, b_i \in \mathbb{R}, 1 \leq i \leq n$ , such that  $\sum_{i=1}^n a_i^2 = \sum_{i=1}^n b_i^2 = 1, \sum_{i=1}^n a_i b_i = 0$ . Prove that

$$\left( \sum_{i=1}^n a_i \right)^2 + \left( \sum_{i=1}^n b_i \right)^2 \leq n.$$

**Example 13 (Macedonia Team Selection Test 2007).** Let  $a, b, c$  be positive real numbers. Prove that

$$1 + \frac{3}{ab + bc + ca} \geq \frac{6}{a + b + c}.$$

**Example 14 (Italian National Olympiad 2007).** a) For each  $n \geq 2$ , find the maximum constant  $c_n$  such that

$$\frac{1}{a_1 + 1} + \frac{1}{a_2 + 1} + \dots + \frac{1}{a_n + 1} \geq c_n,$$

for all positive reals  $a_1, a_2, \dots, a_n$  such that  $a_1 a_2 \cdots a_n = 1$ .

b) For each  $n \geq 2$ , find the maximum constant  $d_n$  such that

$$\frac{1}{2a_1 + 1} + \frac{1}{2a_2 + 1} + \dots + \frac{1}{2a_n + 1} \geq d_n$$

for all positive reals  $a_1, a_2, \dots, a_n$  such that  $a_1 a_2 \cdots a_n = 1$ .

**Example 15 (France Team Selection Test 2007).** Let  $a, b, c, d$  be positive reals such that  $a + b + c + d = 1$ . Prove that

$$6(a^3 + b^3 + c^3 + d^3) \geq a^2 + b^2 + c^2 + d^2 + \frac{1}{8}.$$

**Example 16 (Irish National Mathematical Olympiad 2007).** Suppose  $a, b$  and  $c$  are positive real numbers. Prove that

$$\frac{a + b + c}{3} \leq \sqrt{\frac{a^2 + b^2 + c^2}{3}} \leq \frac{1}{3} \left( \frac{ab}{c} + \frac{bc}{a} + \frac{ca}{b} \right).$$

For each of the inequalities, find conditions on  $a, b$  and  $c$  such that equality holds.

**Example 17 (Vietnam Team Selection Test 2007).** Given a triangle  $ABC$ . Find the minimum of

$$\frac{\cos^2 \frac{A}{2} \cos^2 \frac{B}{2}}{\cos^2 \frac{C}{2}} + \frac{\cos^2 \frac{B}{2} \cos^2 \frac{C}{2}}{\cos^2 \frac{A}{2}} + \frac{\cos^2 \frac{C}{2} \cos^2 \frac{A}{2}}{\cos^2 \frac{B}{2}}.$$

**Example 18 (Greece National Olympiad 2007).** Let  $a, b, c$  be sides of a triangle, show that

$$\frac{(c + a - b)^4}{a(a + b - c)} + \frac{(a + b - c)^4}{b(b + c - a)} + \frac{(b + c - a)^4}{c(c + a - b)} \geq ab + bc + ca.$$

**Example 19 (Bulgaria Team Selection Tests 2007).** Let  $n \geq 2$  is positive integer. Find the best constant  $C(n)$  such that

$$\sum_{i=1}^n x_i \geq C(n) \sum_{1 \leq j < i \leq n} (2x_i x_j + \sqrt{x_i x_j})$$

is true for all real numbers  $x_i \in (0, 1), i = 1, \dots, n$  for which  $(1 - x_i)(1 - x_j) \geq \frac{1}{4}, 1 \leq j < i \leq n$ .

**Example 20 (Poland Second Round 2007).** Let  $a, b, c, d$  be positive real numbers satisfying the following condition:

$$\frac{1}{a} + \frac{1}{b} + \frac{1}{c} + \frac{1}{d} = 4.$$

Prove that:

$$\sqrt[3]{\frac{a^3 + b^3}{2}} + \sqrt[3]{\frac{b^3 + c^3}{2}} + \sqrt[3]{\frac{c^3 + d^3}{2}} + \sqrt[3]{\frac{d^3 + a^3}{2}} \leq 2(a + b + c + d) - 4.$$

**Example 21 (Turkey Team Selection Tests 2007).** Let  $a, b, c$  be positive reals such that their sum is 1. Prove that

$$\frac{1}{ab + 2c^2 + 2c} + \frac{1}{bc + 2a^2 + 2a} + \frac{1}{ac + 2b^2 + 2b} \geq \frac{1}{ab + bc + ac}.$$

**Example 22 (Moldova National Mathematical Olympiad 2007).** Real numbers  $a_1, a_2, \dots, a_n$  satisfy  $a_i \geq \frac{1}{i}$ , for all  $i = \overline{1, n}$ . Prove the inequality

$$(a_1 + 1) \left(a_2 + \frac{1}{2}\right) \cdots \left(a_n + \frac{1}{n}\right) \geq \frac{2^n}{(n+1)!} (1 + a_1 + 2a_2 + \cdots + na_n).$$

**Example 23 (Moldova Team Selection Test 2007).** Let  $a_1, a_2, \dots, a_n \in [0, 1]$ . Denote  $S = a_1^3 + a_2^3 + \cdots + a_n^3$ , prove that

$$\frac{a_1}{2n+1+S-a_1^3} + \frac{a_2}{2n+1+S-a_2^3} + \cdots + \frac{a_n}{2n+1+S-a_n^3} \leq \frac{1}{3}.$$

**Example 24 (Peru Team Selection Test 2007).** Let  $a, b, c$  be positive real numbers, such that

$$a + b + c \geq \frac{1}{a} + \frac{1}{b} + \frac{1}{c}.$$

Prove that

$$a + b + c \geq \frac{3}{a+b+c} + \frac{2}{abc}.$$

**Example 25 (Peru Team Selection Test 2007).** Let  $a, b$  and  $c$  be sides of a triangle. Prove that

$$\frac{\sqrt{b+c-a}}{\sqrt{b}+\sqrt{c}-\sqrt{a}} + \frac{\sqrt{c+a-b}}{\sqrt{c}+\sqrt{a}-\sqrt{b}} + \frac{\sqrt{a+b-c}}{\sqrt{a}+\sqrt{b}-\sqrt{c}} \leq 3.'$$

**Example 26 (Romania Team Selection Tests 2007).** If  $a_1, a_2, \dots, a_n \geq 0$  satisfy  $a_1^2 + \cdots + a_n^2 = 1$ , find the maximum value of the product  $(1 - a_1) \cdots (1 - a_n)$ .

**Example 27 (Romania Team Selection Tests 2007).** Prove that for  $n, p$  integers,  $n \geq 4$  and  $p \geq 4$ , the proposition  $\mathcal{P}(n, p)$

$$\sum_{i=1}^n \frac{1}{x_i^p} \geq \sum_{i=1}^n x_i^p \quad \text{for } x_i \in \mathbb{R}, \quad x_i > 0, \quad i = 1, \dots, n, \quad \sum_{i=1}^n x_i = n,$$

is false.

**Example 28 (Ukraine Mathematical Festival 2007).** Let  $a, b, c$  be positive real numbers and  $abc \geq 1$ . Prove that

(a).

$$\left(a + \frac{1}{a+1}\right) \left(b + \frac{1}{b+1}\right) \left(c + \frac{1}{c+1}\right) \geq \frac{27}{8}.$$

(b).

$$27(a^3+a^2+a+1)(b^3+b^2+b+1)(c^3+c^2+c+1) \geq 64(a^2+a+1)(b^2+b+1)(c^2+c+1).$$

**Example 29 (Asian Pacific Mathematical Olympiad 2007).** Let  $x, y$  and  $z$  be positive real numbers such that  $\sqrt{x} + \sqrt{y} + \sqrt{z} = 1$ . Prove that

$$\frac{x^2 + yz}{\sqrt{2x^2(y+z)}} + \frac{y^2 + zx}{\sqrt{2y^2(z+x)}} + \frac{z^2 + xy}{\sqrt{2z^2(x+y)}} \geq 1.$$



**Example 30 (Brazilian Olympiad Revenge 2007).** Let  $a, b, c \in \mathbb{R}$  with  $abc = 1$ . Prove that

$$a^2 + b^2 + c^2 + \frac{1}{a^2} + \frac{1}{b^2} + \frac{1}{c^2} + 2 \left( a + b + c + \frac{1}{a} + \frac{1}{b} + \frac{1}{c} \right) \geq 6 + 2 \left( \frac{b}{a} + \frac{c}{b} + \frac{a}{c} + \frac{c}{a} + \frac{c}{b} + \frac{b}{c} \right).$$

**Example 31 (India National Mathematical Olympiad 2007).** If  $x, y, z$  are positive real numbers, prove that

$$(x + y + z)^2 (yz + zx + xy)^2 \leq 3(y^2 + yz + z^2)(z^2 + zx + x^2)(x^2 + xy + y^2).$$

**Example 32 (British National Mathematical Olympiad 2007).** Show that for all positive reals  $a, b, c$ ,

$$(a^2 + b^2)^2 \geq (a + b + c)(a + b - c)(b + c - a)(c + a - b).$$

**Example 33 (Korean National Mathematical Olympiad 2007).** For all positive reals  $a, b$ , and  $c$ , what is the value of positive constant  $k$  satisfies the following inequality?

$$\frac{a}{c + kb} + \frac{b}{a + kc} + \frac{c}{b + ka} \geq \frac{1}{2007}.$$

**Example 34 (Hungary-Israel National Mathematical Olympiad 2007).** Let  $a, b, c, d$  be real numbers, such that

$$a^2 \leq 1, a^2 + b^2 \leq 5, a^2 + b^2 + c^2 \leq 14, a^2 + b^2 + c^2 + d^2 \leq 30.$$

Prove that  $a + b + c + d \leq 10$ .

# SOLUTION



Please visit the following links to get the original discussion of the ebook, the problems and solution. We are appreciating every other contribution from you!

<http://www.batdangthuc.net/forum/showthread.php?t=26>

<http://www.batdangthuc.net/forum/showthread.php?t=26&page=2>

<http://www.batdangthuc.net/forum/showthread.php?t=26&page=3>

<http://www.batdangthuc.net/forum/showthread.php?t=26&page=4>

<http://www.batdangthuc.net/forum/showthread.php?t=26&page=5>

<http://www.batdangthuc.net/forum/showthread.php?t=26&page=6>



For Further Reading, Please Review:

- ★ UpComing Vietnam Inequality Forum's Magazine
- ★ Secrets in Inequalities (2 volumes), Pham Kim Hung (hungkhtn)
- ★ Old And New Inequalities, T. Adreescu, V. Cirtoaje, M. Lascu, G. Dospinescu
- ★ Inequalities and Related Issues, Nguyen Van Mau



We thank a lot to Mathlinks Forum and their member for the reference to problems and some nice solutions from them!

**Problem 1 (1, Iran National Mathematical Olympiad 2007).** Assume that  $a, b, c$  are three different positive real numbers. Prove that

$$\left| \frac{a+b}{a-b} + \frac{b+c}{b-c} + \frac{c+a}{c-a} \right| > 1.$$

**Solution 1 (pi3.14).** Due to the symmetry, we can assume  $a > b > c$ . Let  $a = c + x; b = c + y$ , then  $x > y > 0$ . We have

$$\begin{aligned} & \left| \frac{a+b}{a-b} + \frac{b+c}{b-c} + \frac{c+a}{c-a} \right| \\ &= \frac{2c+x+y}{x-y} + \frac{2c+y}{y} - \frac{2c+x}{x} \\ &= 2c \left( \frac{1}{x-y} + \frac{1}{y} - \frac{1}{x} \right) + \frac{x+y}{x-y}. \end{aligned}$$

We have

$$\begin{aligned} 2c \left( \frac{1}{x-y} + \frac{1}{y} - \frac{1}{x} \right) &= 2c \left( \frac{1}{x-y} + \frac{x-y}{xy} \right) > 0. \\ \frac{x+y}{x-y} &> 1. \end{aligned}$$

Thus

$$\left| \frac{a+b}{a-b} + \frac{b+c}{b-c} + \frac{c+a}{c-a} \right| > 1.$$

**Solution 2 (2, Mathlinks, posted by NguyenDungTN).** Let

$$\frac{a+b}{a-b} = x; \frac{b+c}{b-c} = y; \frac{a+c}{c-a} = z;$$

Then

$$xy + yz + xz = 1.$$

By Cauchy-Schwarz Inequality

$$(x+y+z)^2 \geq 3(xy+yz+zx) = 3 \Rightarrow |x+y+z| \geq \sqrt{3} > 1.$$

We are done.

▽

**Problem 2 (2, Iran National Mathematical Olympiad 2007).** Find the largest real  $T$  such that for each non-negative real numbers  $a, b, c, d, e$  such that  $a+b = c+d+e$ , then

$$\sqrt{a^2 + b^2 + c^2 + d^2 + e^2} \geq T(\sqrt{a} + \sqrt{b} + \sqrt{c} + \sqrt{d} + \sqrt{e})^2$$

**Solution 3 (NguyenDungTN).** Let  $a = b = 3, c = d = e = 2$ , we find

$$\frac{\sqrt{30}}{6(\sqrt{3} + \sqrt{2})^2} \geq T.$$

With this value of  $T$ , we will prove the inequality. Indeed, let  $a + b = c + d + e = X$ . By Cauchy-Schwarz Inequality

$$\begin{aligned} a^2 + b^2 &\geq \frac{(a+b)^2}{2} = \frac{X^2}{2}c^2 + d^2 + e^2 \geq \frac{(c+d+e)^2}{3} = \frac{X^2}{3} \\ &\Rightarrow \sqrt{a^2 + b^2 + c^2 + d^2 + e^2} \geq \frac{5X^2}{6} \quad (1) \end{aligned}$$

By Cauchy-Schwarz Inequality, we also have

$$\begin{aligned} \sqrt{a} + \sqrt{b} &\leq \sqrt{2(a+b)} = \sqrt{2X}\sqrt{c} + \sqrt{d} + \sqrt{e} \leq \sqrt{3(c+d+e)} = 3X \\ &\Rightarrow (\sqrt{a} + \sqrt{b} + \sqrt{c} + \sqrt{d} + \sqrt{e})^2 \leq (\sqrt{2} + \sqrt{3})^2 X^2 \quad (2) \end{aligned}$$

From (1) and (2), we have

$$\frac{\sqrt{a^2 + b^2 + c^2 + d^2 + e^2}}{(\sqrt{a} + \sqrt{b} + \sqrt{c} + \sqrt{d} + \sqrt{e})^2} \geq \frac{\sqrt{30}}{6(\sqrt{3} + \sqrt{2})^2}.$$

Equality holds for  $\frac{2a}{3} = \frac{2b}{3} = c = d = e$ .

▽

**Problem 3 (3, Middle European Mathematical Olympiad 2007).** Let  $a, b, c, d$  non-negative such that  $a + b + c + d = 4$ . Prove that

$$a^2bc + b^2cd + c^2da + d^2ab \leq 4.$$

**Solution 4 (mathlinks, reposted by pi3.14).** Let  $\{p, q, r, s\} = \{a, b, c, d\}$  and  $p \geq q \geq r \geq s$ . By rearrangement Inequality, we have

$$\begin{aligned} a^2bc + b^2cd + c^2da + d^2ab &= a(abc) + b(bcd) + c(cda) + d(dab) \\ &\leq p(pqr) + q(pqs) + r(prs) + s(qrs) = (pq + rs)(pr + qs) \\ &\leq \left( \frac{pq + rs + pr + qs}{2} \right)^2 = \frac{1}{4}(p + s)^2(q + r)^2 \\ &\leq \frac{1}{4} \left( \left( \frac{p + q + r + s}{2} \right)^2 \right)^2 = 4. \end{aligned}$$

Equality holds for  $q = r = 1, p + s = 2$ . Easy to refer  $(a, b, c, d) = (1, 1, 1, 1), (2, 1, 1, 0)$  or permutations.

▽

**Problem 4 ( 5- Revised by VanDHHK).** Let  $a, b, c$  be three side-lengths of a triangle such that  $a + b + c = 3$ . Find the minimum of  $a^2 + b^2 + c^2 + \frac{4abc}{3}$

**Solution 5.** Let  $a = x + y, b = y + z, c = z + x$ , we have

$$x + y + z = \frac{3}{2}.$$

Consider

$$\begin{aligned} & a^2 + b^2 + c^2 + \frac{4abc}{3} \\ &= \frac{(a^2 + b^2 + c^2)(a + b + c) + 4abc}{3} \\ &= \frac{2((x + y)^2 + (y + z)^2 + (z + x)^2)(x + y + z) + 4(x + y)(y + z)(z + x)}{3} \\ &= \frac{4(x^3 + y^3 + z^3 + 3x^2y + 3xy^2 + 3y^2z + 3yz^2 + 3z^2x + 3zx^2 + 5xyz)}{3} \\ &= \frac{4((x + y + z)^3 - xyz)}{3} \\ &= \frac{4\left(\frac{26}{27}(x + y + z)^3 + \left(\frac{x+y+z}{3}\right)^3 - xyz\right)}{3} \\ &\geq \frac{4\left(\frac{26}{27}(x + y + z)^3\right)}{3} = \frac{13}{3}. \end{aligned}$$

**Solution 6 (2, DDuLam).** Using the familiar Inequality (equivalent to Schur)

$$abc \geq (b + c - a)(c + a - b)(a + b - c) \Rightarrow abc \geq \frac{4}{3}(ab + bc + ca) - 3.$$

Therefore

$$\begin{aligned} P &\geq a^2 + b^2 + c^2 + \frac{16}{9}(ab + bc + ca) - 4 \\ &= (a + b + c)^2 - \frac{2}{9}(ab + bc + ca) - 4 \geq 5 - \frac{2}{27}(a + b + c)^2 = 4 + \frac{1}{3}. \end{aligned}$$

Equality holds when  $a = b = c = 1$ .

**Solution 7 (3, pi3.14).** With the conventional denotation in triangle, we have

$$abc = 4pRr, \quad a^2 + b^2 + c^2 = 2p^2 - 8Rr - 2r^2.$$

Therefore

$$a^2 + b^2 + c^2 + \frac{4}{3}abc = \frac{9}{2} - 2r^2.$$

Moreover,

$$p \geq 3\sqrt{3}r \Rightarrow r^2 \leq \frac{1}{6}.$$

Thus

$$a^2 + b^2 + c^2 + \frac{4}{3}abc \geq 4\frac{1}{3}.$$

▽

**Problem 5 (7, China Northern Mathematical Olympiad 2007).** Let  $a, b, c$  be positive real numbers such that  $abc = 1$ . Prove that

$$\frac{a^k}{a+b} + \frac{b^k}{b+c} + \frac{c^k}{c+a} \geq \frac{3}{2}.$$

for any positive integer  $k \geq 2$ .

**Solution 8 (Secrets In Inequalities, hungkhtn).** We have

$$\begin{aligned} \frac{a^k}{a+b} + \frac{b^k}{b+c} + \frac{c^k}{c+a} &\geq \frac{3}{2} \\ \Leftrightarrow a^{k-1} + b^{k-1} + c^{k-1} &\geq \frac{3}{2} + \frac{a^{k-1}b}{a+b} + \frac{b^{k-1}c}{b+c} + \frac{c^{k-1}a}{c+a} \end{aligned}$$

By AM-GM Inequality, we have

$$a+b \geq 2\sqrt{ab}, b+c \geq 2\sqrt{bc}, c+a \geq 2\sqrt{ca}.$$

So, it remains to prove that

$$a^{k-\frac{3}{2}}b^{\frac{1}{2}} + b^{k-\frac{3}{2}}c^{\frac{1}{2}} + c^{k-\frac{3}{2}}a^{\frac{1}{2}} + 3 \leq 2(a^{k-1} + b^{k-1} + c^{k-1}).$$

This follows directly by AM-GM inequality, since

$$a^{k-1} + b^{k-1} + c^{k-1} \geq 3\sqrt[3]{a^{k-1}b^{k-1}c^{k-1}} = 3$$

and

$$(2k-3)a^{k-1} + b^{k-1} \geq (2k-2)a^{k-\frac{3}{2}}b^{\frac{1}{2}}$$

$$(2k-3)b^{k-1} + c^{k-1} \geq (2k-2)b^{k-\frac{3}{2}}c^{\frac{1}{2}}$$

$$(2k-3)c^{k-1} + a^{k-1} \geq (2k-2)c^{k-\frac{3}{2}}a^{\frac{1}{2}}$$

Adding up these inequalities, we have the desired result.

▽

**Problem 6 (8, Revised by NguyenDungTN).** Let  $a, b, c > 0$  such that  $a+b+c=1$ . Prove that:

$$\frac{a^2}{b} + \frac{b^2}{c} + \frac{c^2}{a} \geq 3(a^2 + b^2 + c^2).$$

**Solution 9.** By Cauchy-Schwarz Inequality:

$$\frac{a^2}{b} + \frac{b^2}{c} + \frac{c^2}{a} \geq \frac{(a^2 + b^2 + c^2)^2}{a^2b + b^2c + c^2a}.$$

It remains to prove that

$$\begin{aligned} \frac{(a^2 + b^2 + c^2)^2}{a^2b + b^2c + c^2a} &\geq 3(a^2 + b^2 + c^2) \\ \Leftrightarrow (a^2 + b^2 + c^2)(a + b + c) &\geq 3(a^2b + b^2c + c^2a) \\ \Leftrightarrow a^3 + b^3 + c^3 + ab^2 + bc^2 + ca^2 &\geq 2(a^2b + b^2c + c^2a) \\ \Leftrightarrow a(a - b)^2 + b(b - c)^2 + c(c - a)^2 &\geq 0. \end{aligned}$$

So we are done!

**Solution 10 (2, By Zaizai).**

$$\begin{aligned} \frac{a^2}{b} + \frac{b^2}{c} + \frac{c^2}{a} &\geq 3(a^2 + b^2 + c^2) \\ \Leftrightarrow \sum \left( \frac{a^2}{b} - 2a + b \right) &\geq 3(a^2 + b^2 + c^2) - (a + b + c)^2 \\ \Leftrightarrow \sum \frac{(a - b)^2}{b} &\geq (a - b)^2 + (b - c)^2 + (c - a)^2 \\ \Leftrightarrow \sum (a - b)^2 \left( \frac{1}{b} - 1 \right) &\geq 0 \\ \Leftrightarrow \sum \frac{(a - b)^2(a + c)}{b} &\geq 0. \end{aligned}$$

This ends the solution, too.

▽

**Problem 7 (9, Romania Junior Balkan Team Selection Tests 2007).** . Let  $a, b, c$  be three positive reals such that

$$\frac{1}{a + b + 1} + \frac{1}{b + c + 1} + \frac{1}{c + a + 1} \geq 1.$$

Show that

$$a + b + c \geq ab + bc + ca.$$

**Solution 11 (Mathlinks, Reposted by NguyenDungTN).** By Cauchy-Schwarz Inequality, we have

$$(a + b + 1)(a + b + c^2) \geq (a + b + c)^2.$$

Therefore

$$\frac{1}{a+b+1} \leq \frac{c^2+a+b}{(a+b+c)^2},$$

or

$$\begin{aligned} \frac{1}{a+b+1} + \frac{1}{b+c+1} + \frac{1}{c+a+1} &\leq \frac{a^2+b^2+c^2+2(a+b+c)}{(a+b+c)^2} \\ \Rightarrow a^2+b^2+c^2+2(a+b+c) &\geq (a+b+c)^2 \\ \Rightarrow a+b+c &\geq ab+bc+ca. \end{aligned}$$

**Solution 12 (DDucLam).** Assume that  $a+b+c = ab+bc+ca$ , we have to prove that

$$\begin{aligned} \frac{1}{a+b+1} + \frac{1}{b+c+1} + \frac{1}{c+a+1} &\leq 1 \\ \Leftrightarrow \frac{a+b}{a+b+1} + \frac{b+c}{b+c+1} + \frac{c+a}{c+a+1} &\geq 2 \end{aligned}$$

By Cauchy-Schwarz Inequality,

$$\text{LHS} \geq \frac{(a+b+b+c+c+a)^2}{\sum_{cyc}(a+b)(a+b+1)} = 2.$$

We are done

**Comment.** This second very beautiful solution uses Contradiction method. If you can't understand the principal of this method, have a look at *Sang Tao Bat Dang Thuc*, or *Secrets In Inequalities*, written by Pham Kim Hung.

▽

**Problem 8 (10, Romanian JBTST V 2007).** Let  $x, y, z$  be non-negative real numbers.

Prove that

$$\frac{x^3+y^3+z^3}{3} \geq xyz + \frac{3}{4}|(x-y)(y-z)(z-x)|.$$

**Solution 13 (vandhkh).** We have

$$\begin{aligned} \frac{x^3+y^3+z^3}{3} &\geq xyz + \frac{3}{4}|(x-y)(y-z)(z-x)| \\ \Leftrightarrow \frac{x^3+y^3+z^3}{3} - xyz &\geq \frac{3}{4}|(x-y)(y-z)(z-x)| \\ \Leftrightarrow ((x-y)^2 + (y-z)^2 + (z-x)^2)((x+y) + (y+z) + (z+x)) &\geq 9|(x-y)(y-z)(z-x)|. \end{aligned}$$

Notice that

$$x+y \geq |x-y|; y+z \geq |y-z|; z+x \geq |z-x|,$$

and by AM-GM Inequality,

$$((x-y)^2 + (y-z)^2 + (z-x)^2)(|x-y| + |y-z| + |z-x|) \geq 9|(x-y)(y-z)(z-x)|.$$

So we are done. Equality holds for  $x = y = z$ .



**Solution 14 (Secrets In Inequalities, hungkhtn).** The inequality is equivalent to

$$(x + y + z) \sum (x - y)^2 \geq \frac{9}{2} |(x - y)(y - z)(z - x)|.$$

By the entirely mixing variable method, it is enough to prove when  $z = 0$

$$x^3 + y^3 \geq \frac{9}{4} |xy(x - y)|.$$

This last inequality can be checked easily.

▽

**Problem 9 (11, Yugoslavia National Olympiad 2007).** Let  $k$  be a given natural number. Prove that for any positive numbers  $x, y, z$  with the sum 1, the following inequality holds

$$\frac{x^{k+2}}{x^{k+1} + y^k + z^k} + \frac{y^{k+2}}{y^{k+1} + z^k + x^k} + \frac{z^{k+2}}{z^{k+1} + x^k + y^k} \geq \frac{1}{7}.$$

When does equality occur?

**Solution 15 (NguyenDungTN).** We can assume that  $x \geq y \geq z$ . By this assumption, easy to refer that

$$\begin{aligned} \frac{x^{k+1}}{x^{k+1} + y^k + z^k} &\geq \frac{y^{k+1}}{y^{k+1} + z^k + x^k} \geq \frac{z^{k+1}}{z^{k+1} + x^k + y^k}; \\ z^{k+1} + y^k + x^k &\geq y^{k+1} + x^k + z^k \geq x^{k+1} + z^k + y^k; \end{aligned}$$

and

$$x^k \geq y^k \geq z^k.$$

By Chebyshev Inequality, we have

$$\begin{aligned} &\frac{x^{k+2}}{x^{k+1} + y^k + z^k} + \frac{y^{k+2}}{y^{k+1} + z^k + x^k} + \frac{z^{k+2}}{z^{k+1} + x^k + y^k} \\ &\geq \frac{x + y + z}{3} \left( \frac{x^{k+1}}{x^{k+1} + y^k + z^k} + \frac{y^{k+1}}{y^{k+1} + z^k + x^k} + \frac{z^{k+1}}{z^{k+1} + x^k + y^k} \right) \\ &= \frac{1}{3} \left( \frac{x^{k+1}}{x^{k+1} + y^k + z^k} + \frac{y^{k+1}}{y^{k+1} + z^k + x^k} + \frac{z^{k+1}}{z^{k+1} + x^k + y^k} \right) \frac{\sum_{cyc} (x^{k+1} + y^k + z^k)}{\sum_{cyc} (x^{k+1} + y^k + z^k)} \\ &= \frac{1}{3} \left( \sum_{cyc} \left( \frac{x^{k+1}}{x^{k+1} + y^k + z^k} \sum_{cyc} (x^{k+1} + y^k + z^k) \frac{1}{\sum_{cyc} (x^{k+1} + y^k + z^k)} \right) \right) \\ &\geq \frac{1}{3} (x^{k+1} + y^{k+1} + z^{k+1}) \cdot \frac{1}{\sum_{cyc} (x^{k+1} + y^k + z^k)} = \frac{x^{k+1} + y^{k+1} + z^{k+1}}{x^{k+1} + y^{k+1} + z^{k+1} + 2(x^k + y^k + z^k)} \end{aligned}$$

Also by Chebyshev Inequality,

$$3(x^{k+1} + y^{k+1} + z^{k+1}) \geq 3 \frac{x+y+z}{3} (x^k + y^k + z^k) = x^k + y^k + z^k.$$

Thus

$$\frac{x^{k+1} + y^{k+1} + z^{k+1}}{x^{k+1} + y^{k+1} + z^{k+1} + 2(x^k + y^k + z^k)} \geq \frac{x^{k+1} + y^{k+1} + z^{k+1}}{x^{k+1} + y^{k+1} + z^{k+1} + 6(x^k + y^k + z^k)} = \frac{1}{7}.$$

So we are done. Equality holds for  $a = b = c = \frac{1}{3}$ .

▽

**Problem 10 (Macedonia Team Selection Test 2007).** Let  $a, b, c$  be positive real numbers.

Prove that

$$1 + \frac{3}{ab + bc + ca} \geq \frac{6}{a + b + c}.$$

**Solution 16 (VoDanh).** The inequality is equivalent to

$$a + b + c + \frac{3(a + b + c)}{ab + bc + ca} \geq 6.$$

By AM-GM Inequality,

$$a + b + c + \frac{3(a + b + c)}{ab + bc + ca} \geq 2\sqrt{\frac{3(a + b + c)^2}{ab + bc + ca}}.$$

It is obvious that  $(a + b + c)^2 \geq 3(ab + bc + ca)$ , so we are done!

▽

**Problem 11 (14, Italian National Olympiad 2007).** a). For each  $n \geq 2$ , find the maximum constant  $c_n$  such that:

$$\frac{1}{a_1 + 1} + \frac{1}{a_2 + 1} + \dots + \frac{1}{a_n + 1} \geq c_n,$$

for all positive reals  $a_1, a_2, \dots, a_n$  such that  $a_1 a_2 \cdots a_n = 1$ .

▽

b). For each  $n \geq 2$ , find the maximum constant  $d_n$  such that

$$\frac{1}{2a_1 + 1} + \frac{1}{2a_2 + 1} + \dots + \frac{1}{2a_n + 1} \geq d_n,$$

for all positive reals  $a_1, a_2, \dots, a_n$  such that  $a_1 a_2 \cdots a_n = 1$ .

**Solution 17 (Mathlinks, reposted by NguyenDungTN).** a). Let

$$a_1 = \epsilon^{n-1}, a_k = \frac{1}{\epsilon} \forall k \neq 1,$$

then let  $\epsilon \rightarrow 0$ , we easily get  $c_n \leq 1$ . We will prove the inequality with this value of  $c_n$ . Without loss of generality, assume that  $a_1 \leq a_2 \leq \dots \leq a_n$ . Since  $a_1 a_2 \leq 1$ , we have

$$\sum_{k=1}^n \frac{1}{a_k + 1} \geq \frac{1}{a_1 + 1} + \frac{1}{a_2 + 1} = \frac{1}{a_1 + 1} + \frac{a_1}{a_2 + a_1 a_2} \geq \frac{1}{a_1 + 1} + \frac{a_1}{a_1 + 1} = 1.$$

This ends the proof.

b). Consider  $n = 2$ , it is easy to get  $d_2 = \frac{2}{3}$ . Indeed, let  $a_1 = a, a_2 = \frac{1}{a}$ . The inequality becomes

$$\begin{aligned} \frac{1}{2a+1} + \frac{a}{a+2} \geq \frac{2}{3} &\Leftrightarrow 3(a+2) + 3a(2a+1) \geq 2(2a+1)(a+2) \\ &\Leftrightarrow (a-1)^2 \geq 0. \end{aligned}$$

When  $n \geq 3$ , similar to (a), we will show that  $d_n = 1$ . Indeed, without loss of generality, we may assume that

$$a_1 \leq a_2 \leq \dots \leq a_n \Rightarrow a_1 a_2 a_3 \leq 1.$$

Let

$$x = \sqrt[9]{\frac{a_2 a_3}{a_1^2}}, y = \sqrt[9]{\frac{a_1 a_3}{a_2^2}}, z = \sqrt[9]{\frac{a_1 a_2}{a_3^2}}$$

then  $a_1 \leq \frac{1}{x^3}, a_2 \leq \frac{1}{y^3}, a_3 \leq \frac{1}{z^3}, xyz = 1$ . Thus

$$\begin{aligned} \sum_{k=1}^n \frac{1}{a_k + 1} &\geq \sum_{k=1}^3 \frac{1}{a_k + 1} \geq \frac{x^3}{x^3 + 2} + \frac{y^3}{y^3 + 2} + \frac{z^3}{z^3 + 2} \\ &= \frac{x^2}{x^2 + 2yz} + \frac{y^2}{y^2 + 2xz} + \frac{z^2}{z^2 + 2xy} \\ &\geq \frac{x^2}{x^2 + y^2 + z^2} + \frac{y^2}{x^2 + y^2 + z^2} + \frac{z^2}{x^2 + y^2 + z^2} = 1. \end{aligned}$$

This ends the proof.

▽

**Problem 12 (15, France Team Selection Test 2007).** . Let  $a, b, c, d$  be positive reals such that  $a + b + c + d = 1$ . Prove that:

$$6(a^3 + b^3 + c^3 + d^3) \geq a^2 + b^2 + c^2 + d^2 + \frac{1}{8}.$$

**Solution 18 (NguyenDungTN).** By AM-GM Inequality

$$2a^3 + \frac{1}{4^3} \geq \frac{3a^2}{4}a^2 + \frac{1}{4^2} \geq \frac{a}{2}.$$

Therefore

$$6(a^3 + b^3 + c^3 + d^3) + \frac{3}{16} \geq \frac{9(a^2 + b^2 + c^2 + d^2)}{4}$$

$$\frac{5(a^2 + b^2 + c^2 + d^2)}{4} + \frac{5}{16} \geq \frac{5(a + b + c + d)}{8} = \frac{5}{8}$$

Adding up two of them, we get

$$6(a^3 + b^3 + c^3 + d^3) \geq a^2 + b^2 + c^2 + d^2 + \frac{1}{8}$$

**Solution 19 (Zaizai).** We know that

$$6a^3 \geq a^2 + \frac{5a}{8} - \frac{1}{8} \Leftrightarrow \frac{(4a-1)^2(3a+1)}{8} \geq 0$$

Adding up four similar inequalities, we are done!

▽

**Problem 13 (16, Revised by NguyenDungTN).** Suppose  $a, b$  and  $c$  are positive real numbers. Prove that

$$\frac{a+b+c}{3} \leq \sqrt{\frac{a^2+b^2+c^2}{3}} \leq \frac{1}{3} \left( \frac{bc}{a} + \frac{ca}{b} + \frac{ab}{c} \right).$$

**Solution 20.** The left-hand inequality is just Cauchy-Schwarz Inequality. We will prove the right one. Let

$$\frac{bc}{a} = x, \frac{ca}{b} = y, \frac{ab}{c} = z.$$

The inequality becomes

$$\sqrt{\frac{xy+yz+zx}{3}} \leq \frac{x+y+z}{3}.$$

Squaring both sides, the inequality becomes

$$(x+y+z)^2 \geq 3(xy+yz+zx) \Leftrightarrow (x-y)^2 + (y-z)^2 + (z-x)^2 \geq 0,$$

which is obviously true.

▽

**Problem 14 (17, Vietnam Team Selection Test 2007).** Given a triangle  $ABC$ . Find the minimum of:

$$\sum \frac{\cos^2(\frac{A}{2})\cos^2(\frac{B}{2})}{\cos^2(\frac{C}{2})}$$

**Solution 21 (pi3.14).** We have

$$\begin{aligned} T &= \sum \frac{\cos^2(\frac{A}{2})\cos^2(\frac{B}{2})}{\cos^2(\frac{C}{2})} \\ &= \sum \frac{(1 + \cos A)(1 + \cos B)}{2(1 + \cos C)}. \end{aligned}$$

Let  $a = \tan\frac{A}{2}$ ;  $b = \tan\frac{B}{2}$ ;  $c = \tan\frac{C}{2}$ . We have  $ab + bc + ca = 1$ . So

$$\begin{aligned} T &= \sum \frac{(1 + a^2)}{(1 + b^2)(1 + c^2)} = \sum \frac{1}{\frac{(1+b^2)(1+c^2)}{1+a^2}} \\ &= \sum \frac{1}{\frac{(ab+bc+ca+b^2)(ab+bc+ca+c^2)}{(ab+bc+ca+a^2)}} \\ &= \sum \frac{1}{\frac{(a+b)(c+b)(a+c)(b+c)}{(b+a)(b+c)}} \\ &= \sum \frac{1}{(b+c)^2} \end{aligned}$$

By Iran96 Inequality, we have

$$\frac{1}{(b+c)^2} + \frac{1}{(c+a)^2} + \frac{1}{(a+b)^2} \geq \frac{9}{4(ab+bc+ca)}.$$

Thus  $F \geq \frac{9}{4}$  Equality holds when ABC is equilateral.

▽

**Problem 15 (18, Greece National Olympiad 2007).** . Let  $a, b, c$  be sides of a triangle, show that

$$\frac{(b+c-a)^4}{a(a+b-c)} + \frac{(c+a-b)^4}{b(b+c-a)} + \frac{(b+c-a)^4}{a(c+a-b)} \geq ab+bc+ca.$$

**Solution 22 (NguyenDungTN).** Since  $a, b, c$  are three sides of a triangle, we can substitute

$$a = y + z, b = z + x, c = x + y.$$

The inequality becomes

$$\frac{8x^4}{(x+y)y} + \frac{8y^4}{(y+z)z} + \frac{8z^4}{(z+x)x} \geq x^2 + y^2 + z^2 + 3(xy + yz + zx).$$

By Cauchy-Schwarz Inequality, we have

$$\frac{8x^4}{(x+y)y} + \frac{8y^4}{(y+z)z} + \frac{8z^4}{(z+x)x} \geq \frac{8(x^2 + y^2 + z^2)^2}{x^2 + y^2 + z^2 + xy + yz + zx}.$$

We will prove that

$$\begin{aligned} & \frac{8(x^2 + y^2 + z^2)^2}{x^2 + y^2 + z^2 + xy + yz + zx} \geq x^2 + y^2 + z^2 + 3(xy + yz + zx) \\ \Leftrightarrow & 8(x^2 + y^2 + z^2)^2 \geq (x^2 + y^2 + z^2 + xy + yz + zx)(x^2 + y^2 + z^2 + 3(xy + yz + zx)) \\ \Leftrightarrow & 8 \sum x^4 + 16 \sum x^2 y^2 \geq \sum x^4 + 2 \sum x^2 y^2 + \\ & + 4 \sum x^3(y + z) + 12xyz(x + y + z) + 3 \sum x^2 y^2 + 6xyz(x + y + z) \\ \Leftrightarrow & 7 \sum x^4 + 11 \sum x^2 y^2 \geq 4 \sum x^3(y + z) + 10xyz(x + y + z). \end{aligned}$$

By AM-GM and Schur Inequality

$$\begin{aligned} & 3 \sum x^4 + 11 \sum x^2 y^2 \geq 14xyz(x + y + z); \\ & 4 \left( \sum x^4 + xyz(x + y + z) \right) \geq 4 \sum x^3(y + z) \end{aligned}$$

Adding up two inequalities, we are done!

**Solution 23 (2, DducLam).** By AM-GM Inequality, we have

$$\frac{(b + c - a)^4}{a(a + b - c)} + a(a + b - c) \geq 2(b + c - a)^2.$$

Construct two similar inequalities, then adding up, we have

$$\begin{aligned} & \frac{(b + c - a)^4}{a(a + b - c)} + \frac{(c + a - b)^4}{b(b + c - a)} + \frac{(b + c - a)^4}{a(c + a - b)} \\ & \geq 2[3(a^2 + b^2 + c^2) - 2(ab + bc + ca)] - (a^2 + b^2 + c^2) \\ & = 5(a^2 + b^2 + c^2) - 4(ab + bc + ca) \geq ab + bc + ca. \end{aligned}$$

We are done!

▽

**Problem 16 (20, Poland Second Round 2007).** . Let  $a, b, c, d$  be positive real numbers satisfying the following condition  $\frac{1}{a} + \frac{1}{b} + \frac{1}{c} + \frac{1}{d} = 4$  Prove that:

$$\sqrt[3]{\frac{a^3 + b^3}{2}} + \sqrt[3]{\frac{b^3 + c^3}{2}} + \sqrt[3]{\frac{c^3 + d^3}{2}} + \sqrt[3]{\frac{d^3 + a^3}{2}} \leq 2(a + b + c + d) - 4.$$

**Solution 24 (Mathlinks, reposted by NguyenDungTN).** First, we show that

$$\sqrt[3]{\frac{a^3 + b^3}{2}} \leq \frac{a^2 + b^2}{a + b},$$

which is equivalent to

$$(a - b)^4(a^2 + ab + b^2) \geq 0.$$

Therefore, we refer that

$$\sqrt[3]{\frac{a^3 + b^3}{2}} + \sqrt[3]{\frac{b^3 + c^3}{2}} + \sqrt[3]{\frac{c^3 + d^3}{2}} + \sqrt[3]{\frac{d^3 + a^3}{2}} \leq \frac{a^2 + b^2}{a + b} + \frac{b^2 + c^2}{b + c} + \frac{c^2 + d^2}{c + d} + \frac{d^2 + a^2}{d + a}$$

It remains to prove that

$$\frac{a^2 + b^2}{a + b} + \frac{b^2 + c^2}{b + c} + \frac{c^2 + d^2}{c + d} + \frac{d^2 + a^2}{d + a} \leq 2(a + b + c + d) - 4.$$

However,

$$a + b - \frac{a^2 + b^2}{a + b} = \frac{2ab}{a + b} = \frac{2}{\frac{1}{a} + \frac{1}{b}},$$

So, due to Cauchy-Schwarz Inequality, we get

$$\begin{aligned} & 2(a + b + c + d) - \left( \frac{a^2 + b^2}{a + b} + \frac{b^2 + c^2}{b + c} + \frac{c^2 + d^2}{c + d} + \frac{d^2 + a^2}{d + a} \right) \\ &= 2 \sum \frac{1}{\frac{1}{a} + \frac{1}{b}} \geq 2 \frac{4^2}{2\left(\frac{1}{a} + \frac{1}{b} + \frac{1}{c} + \frac{1}{d}\right)} = \frac{32}{8} = 4 \end{aligned}$$

This ends the proof.

▽

**Problem 17 (21, Turkey Team Selection Tests 2007).** . Let  $a, b, c$  be positive reals such that their sum is 1. Prove that:

$$\frac{1}{ab + 2c^2 + 2c} + \frac{1}{bc + 2a^2 + 2a} + \frac{1}{ac + 2b^2 + 2b} \geq \frac{1}{ab + bc + ac}.$$

**Solution 25 (NguyenDungTN).** First, we will prove that

$$\frac{ab + ac + bc}{ab + 2c^2 + 2c} \geq \frac{ab}{ab + ac + bc}.$$

Indeed, this is equivalent to

$$a^2b^2 + b^2c^2 + c^2a^2 + 2abc(a + b + c) \geq a^2b^2 + 2abc^2 + 2abc,$$

which is always true since  $2abc(a + b + c) = 2abc$  and due to AM-GM Inequality

$$a^2c^2 + b^2c^2 \geq 2abc^2.$$

Similarly, we have

$$\frac{ab + ac + bc}{bc + 2a^2 + 2a} \geq \frac{bc}{ab + ac + bc}.$$

$$\frac{ab + ac + bc}{ac + 2b^2 + 2b} \geq \frac{ca}{ab + ac + bc}.$$

Adding up three inequalities, we are done!

▽

**Problem 18 (22, Moldova National Mathematical Olympiad 2007).** Real numbers  $a_1, a_2, \dots, a_n$  satisfy  $a_i \geq \frac{1}{i}$ , for all  $i = \overline{1, n}$ . Prove the inequality

$$(a_1 + 1) \left(a_2 + \frac{1}{2}\right) \cdots \left(a_n + \frac{1}{n}\right) \geq \frac{2^n}{(n+1)!} (1 + a_1 + 2a_2 + \cdots + na_n).$$

**Solution 26 (NguyenDungTN).** This inequality is equivalent to

$$(a_1 + 1)(2a_2 + 1) \cdots (na_n + 1) \geq \frac{2^n}{n+1} (1 + a_1 + 2a_2 + \cdots + na_n).$$

It is clearly true when  $n = 1$ . Assume that it is true for  $n = k$ , we have to prove it for  $n = k + 1$ . Indeed,

$$(a_1 + 1)(2a_2 + 1) \cdots (ka_k + 1)((k+1)a_{k+1} + 1) \geq \frac{2^k}{k+1} (1 + a_1 + 2a_2 + \cdots + ka_k)((k+1)a_{k+1} + 1)$$

Let

$$a = (k+1)a_{k+1}s = a_1 + 2a_2 + \cdots + ka_k \Rightarrow s \geq k.$$

We need to show that

$$\frac{2^k}{k+1} (1+s)(1+a) \geq \frac{2^{k+1}}{k+2} (1+s+a)$$

$$\Leftrightarrow 2(as - k) + k(a-1)(s-1) \geq 0.$$

Since  $a \geq 1 \forall k$ , the above one is true for  $n = k + 1$ . The proof ends! Equality holds for  $a_i = \frac{1}{i}, i = \overline{1, n}$ .

**Solution 27 (NguyenDungTN).** The inequality is equivalent to

$$\left(\frac{1+a_1}{2}\right) \left(\frac{1+2a_2}{2}\right) \cdots \left(\frac{1+na_n}{2}\right) \geq \frac{1+a_1+2a_2+\cdots+na_n}{n+1}.$$

Let  $x_i = \frac{ia_i-1}{2} \geq 0$ , it becomes

$$(1+x_1)(1+x_2)\cdots(1+x_n) \geq 1 + \frac{2}{n+1}(x_1+x_2+\cdots+x_n).$$

But

$$(1+x_1)(1+x_2)\cdots(1+x_n) \geq 1 + x_1 + x_2 + \cdots + x_n \geq 1 + \frac{2}{n+1}(x_1+x_2+\cdots+x_n).$$

So we have the desired result.



▽

**Problem 19 (23, Moldova Team Selection Test 2007).** Let  $a_1, a_2, \dots, a_n \in [0, 1]$ . Denote  $S = a_1^3 + a_2^3 + \dots + a_n^3$ . Prove that

$$\frac{a_1}{2n+1+S-a_1^3} + \frac{a_2}{2n+1+S-a_2^3} + \dots + \frac{a_n}{2n+1+S-a_n^3} \leq \frac{1}{3}.$$

**Solution 28 (NguyenDungTN).** By AM-GM Inequality, we have

$$S - a_1^3 + 2(n-1) = (a_2^3 + 2) + (a_3^3 + 2) + \dots + (a_n^3 + 2) \geq 3(a_2 + a_3 + \dots + a_n).$$

Thus

$$\frac{a_1}{2n+1+S-a_1^3} \leq \frac{a_1}{3(1+a_1+a_2+\dots+a_n)} \leq \frac{a_1}{3(a_1+a_2+\dots+a_n)}.$$

Similar for  $a_2, a_3, \dots, a_n$ , we have

$$\begin{aligned} \frac{a_1}{2n+1+S-a_1^3} + \frac{a_2}{2n+1+S-a_2^3} + \dots + \frac{a_n}{2n+1+S-a_n^3} \\ \leq \frac{1}{3} \cdot \frac{a_1+a_2+\dots+a_n}{a_1+a_2+\dots+a_n} = \frac{1}{3}. \end{aligned}$$

The equality holds for  $a_1 = a_2 = \dots = a_n = 1$ .

▽

**Problem 20 (24, Peru Team Selection Test 2007).** Let  $a, b, c$  be positive real numbers, such that

$$a + b + c \geq \frac{1}{a} + \frac{1}{b} + \frac{1}{c}.$$

Prove that:

$$a + b + c \geq \frac{3}{a+b+c} + \frac{2}{abc}.$$

**Solution 29 (NguyenDungTN).** By Cauchy-Schwarz Inequality, we have

$$a + b + c \geq \frac{1}{a} + \frac{1}{b} + \frac{1}{c} \geq \frac{9}{a+b+c} \Rightarrow a + b + c \geq 3.$$

Our inequality is equivalent to

$$(a + b + c)^2 \geq 3 + 2 \left( \frac{1}{ab} + \frac{1}{bc} + \frac{1}{ca} \right).$$

By AM-GM Inequality

$$2 \left( \frac{1}{ab} + \frac{1}{bc} + \frac{1}{ca} \right) \leq \frac{2}{3} \left( \frac{1}{a} + \frac{1}{b} + \frac{1}{c} \right)^2 \leq \frac{2}{3} (a + b + c)^2$$

So it is enough to prove that

$$(a + b + c)^2 \geq 3 + \frac{2}{3}(a + b + c)^2 \Leftrightarrow (a + b + c)^2 \geq 9.$$

This inequality is true due to  $a + b + c \geq 3$ .

**Solution 30 (2, DducLam).** We have

$$a + b + c \geq \frac{2}{3}(a + b + c) + \frac{1}{3}\left(\frac{1}{a} + \frac{1}{b} + \frac{1}{c}\right) \geq \frac{2}{3}(a + b + c) + \frac{3}{a + b + c}.$$

We only need to prove that

$$a + b + c \geq \frac{3}{abc},$$

but this inequality is always true since

$$(a + b + c)^2 \geq \left(\frac{1}{a} + \frac{1}{b} + \frac{1}{c}\right)^2 \geq 3\left(\frac{1}{ab} + \frac{1}{bc} + \frac{1}{ca}\right) = \frac{3}{abc}(a + b + c).$$

▽

**Problem 21 (25, Revised by NguyenDungTN).** Let  $a, b$  and  $c$  be sides of a triangle.

Prove that

$$\frac{\sqrt{b+c-a}}{\sqrt{b}+\sqrt{c}-\sqrt{a}} + \frac{\sqrt{c+a-b}}{\sqrt{c}+\sqrt{a}-\sqrt{b}} + \frac{\sqrt{a+b-c}}{\sqrt{a}+\sqrt{b}-\sqrt{c}} \leq 3.$$

**Solution 31.** Let

$$x = \sqrt{b} + \sqrt{c} - \sqrt{a}, y = \sqrt{c} + \sqrt{a} - \sqrt{b}, z = \sqrt{a} + \sqrt{b} - \sqrt{c},$$

then

$$b + c - a = x^2 - \frac{(x-y)(x-z)}{2}$$

By AM-GM inequality, we have

$$\frac{\sqrt{b+c-a}}{\sqrt{b}+\sqrt{c}-\sqrt{a}} = \sqrt{1 - \frac{(x-y)(x-z)}{2x^2}} \leq 1 - \frac{(x-y)(x-z)}{4x^2}$$

We will prove that

$$x^{-2}(x-y)(x-z) + y^{-2}(y-z)(y-x) + z^{-2}(z-x)(z-y) \geq 0.$$

But this immediately follows the general Schur inequality, with the assumption that

$$x \geq y \geq z \Rightarrow x^{-2} \leq y^{-2} \leq z^{-2}.$$

We are done!

▽

**Problem 22 (26, Romania Team Selection Tests 2007).** If  $a_1, a_2, \dots, a_n \geq 0$  are such that  $a_1^2 + \dots + a_n^2 = 1$ , find the maximum value of the product  $(1 - a_1) \cdots (1 - a_n)$ .

**Solution 32 (hungkhtn, reposted by NguyenDungTN).** We use contradiction method. Assume that  $x_1, x_2, \dots, x_n \in [0, 1]$  such that  $x_1 x_2 \dots x_n = (1 - \frac{1}{\sqrt{2}})^2$ . We will prove

$$f(x_1, x_2, \dots, x_n) = (1 - x_1)^2 + (1 - x_2)^2 + \dots + (1 - x_n)^2 \leq 1 \quad (1)$$

Indeed, first, we prove that:

Lemma: If  $x, y \in [0, 1]$ ,  $x + y + xy \geq 1$  then

$$(1 - x)^2 + (1 - y)^2 \leq (1 - xy)^2.$$

Proof. Notice that

$$\begin{aligned} (1 - x)^2 + (1 - y)^2 - (1 - xy)^2 &= (x + y - 1)^2 - x^2 y^2 \\ &= (x - 1)(y - 1)(x + y + xy - 1) \leq 0. \end{aligned}$$

The lemma is asserted. Return to the problem, let  $k = 1 - \frac{1}{\sqrt{2}}$ . Assume that  $x_1 \leq x_2 \leq \dots \leq x_n$ , then

$$x_1 x_2 x_3 \geq k^2 \Rightarrow x_2 x_3 \geq k^{4/3},$$

thus

$$x_2 + x_3 + x_2 x_3 \geq 2k^{2/3} + k^{4/3} = 1.07 \geq 1.$$

Similarly, we have

$$\begin{aligned} f(x_1, x_2, \dots, x_n) &\leq f(x_1, x_2 x_3, 1, x_4, \dots, x_n) \\ &\leq f(x_1, x_2 x_3 x_4, 1, 1, x_5, \dots, x_n) \leq \dots \leq f(x_1, x_2 x_3 \dots x_n, 1, 1, \dots, 1), \end{aligned}$$

From this, easy to get the final result.

▽

**Problem 23 (28, Ukraine Mathematic Festival 2007).** Let  $a, b, c > 0$  và  $abc \geq 1$ . Prove that

$$\begin{aligned} a). & \left(a + \frac{1}{a+1}\right) \left(b + \frac{1}{b+1}\right) \left(c + \frac{1}{c+1}\right) \geq \frac{27}{8}. \\ b). & 27(a^3 + a^2 + a + 1)(b^3 + b^2 + b + 1)(c^3 + c^2 + c + 1) \\ & \geq 64(a^2 + a + 1)(b^2 + b + 1)(c^2 + c + 1). \end{aligned}$$

**Solution 33 (pi3.14).** Consider the case  $abc = 1$ . Let  $a = \frac{x}{y}$ ,  $b = \frac{y}{z}$ ,  $c = \frac{z}{x}$ . The inequality becomes

$$\sum \frac{\frac{x^2}{y^2} + \frac{x}{y} + 1}{\frac{x}{y} + 1} \geq \frac{27}{8}$$

or

$$8(x^2 + xy + y^2)(y^2 + yz + z^2)(x^2 + zx + z^2) \geq 27xyz(x + y)(y + z)(z + x) \quad (1)$$

We have

$$2(x^2 + xy + y^2) \geq 3\sqrt{xy}(x + y),$$

since

$$2(x^2 + xy + y^2) \geq \frac{3}{2}(x^2 + 2xy + y^2) \geq 3\sqrt{xy}(x + y).$$

Write two similar inequalities, then multiply all of them, we get (1) immediately.

If  $abc > 1$ , we let  $a = ka'$ ;  $b = kb'$ ;  $c = kc'$ ; with  $k = \sqrt[3]{abc}$ . We have  $k > 1$  and  $a'b'c' = 1$ . Then

$$\frac{a^2 + a + 1}{a + 1} \geq \frac{a'^2 + a' + 1}{a' + 1}.$$

Since the inequality is proved for  $a', b', c'$ , this is true for  $a, b, c$  immediately.

b). By AM-GM inequality

$$a^2 + 2 \geq 2a \Rightarrow (a^2 + 1) \geq \frac{2}{3}(a^2 + a + 1).$$

Therefore

$$3(a^3 + a^2 + a + 1) = 3(a + 1)(a^2 + 1) \geq 6\sqrt{a} \cdot \frac{2}{3}(a^2 + a + 1) = 4\sqrt{a}(a^2 + a + 1).$$

Constructing similar inequalities, then multiply all of them, we get

$$27(a^3 + a^2 + a + 1)(b^3 + b^2 + b + 1)(c^3 + c^2 + c + 1) \geq 64(a^2 + a + 1)(b^2 + b + 1)(c^2 + c + 1).$$

**Solution 34 (2, NguyenDungTN).** By AM-GM inequality

$$\frac{a + 1}{4} + \frac{1}{a + 1} \geq 1;$$

$$\frac{3a}{4} + \frac{3}{4} \geq \frac{3}{2}\sqrt{a};$$

Adding up two inequalities, we get

$$a + \frac{1}{a + 1} \geq \frac{3}{2}\sqrt{a}.$$

Similar for  $b, c$ , and finally we have

$$\left(a + \frac{1}{a + 1}\right) \left(b + \frac{1}{b + 1}\right) \left(c + \frac{1}{c + 1}\right) \geq \frac{27}{8}\sqrt{abc} \geq \frac{27}{8}.$$

Equality holds for  $a = b = c = 1$ .

▽

**Problem 24 (29, Asian Pacific Mathematical Olympiad 2007).** Let  $x, y$  and  $z$  be positive real numbers such that  $\sqrt{x} + \sqrt{y} + \sqrt{z} = 1$ . Prove that

$$\frac{x^2 + yz}{\sqrt{2x^2(y+z)}} + \frac{y^2 + zx}{\sqrt{2y^2(z+x)}} + \frac{z^2 + xy}{\sqrt{2z^2(x+y)}} \geq 1.$$

**Solution 35 (NguyenDungTN).** We have the transformation

$$\sum_{cyc} \frac{x^2 + yz}{\sqrt{2x^2(y+z)}} = \sum_{cyc} \frac{(x-y)(x-z)}{\sqrt{2x^2(y+z)}} + \sum_{cyc} \sqrt{\frac{y+z}{2}}.$$

Moreover, by Cauchy-Schwarz Inequality

$$\sum_{cyc} \sqrt{\frac{y+z}{2}} \geq \sum_{cyc} \frac{\sqrt{y} + \sqrt{z}}{2} = 1.$$

So it is enough to prove that

$$\sum_{cyc} \frac{(x-y)(x-z)}{\sqrt{2x^2(y+z)}} \geq 0$$

Without loss of generality, assume that  $x \geq y \geq z$ , then

$$\frac{1}{\sqrt{2x^2(y+z)}} \leq \frac{1}{\sqrt{2y^2(z+x)}} \leq \frac{1}{\sqrt{2z^2(x+y)}}.$$

Using the general Schur Inequality, we have the desired result.

▽

**Problem 25 (30, Brazilian Olympiad Revenge 2007).** Let  $a, b, c \in \mathbb{R}$  with  $abc = 1$ . Prove that

$$a^2 + b^2 + c^2 + \frac{1}{a^2} + \frac{1}{b^2} + \frac{1}{c^2} + 2 \left( a + b + c + \frac{1}{a} + \frac{1}{b} + \frac{1}{c} \right) \geq 6 + 2 \left( \frac{a}{b} + \frac{b}{c} + \frac{c}{a} + \frac{b}{a} + \frac{c}{b} + \frac{a}{c} \right).$$

**Solution 36 (NguyenDungTN).** Since  $abc = 1$ , we have

$$a^2 + b^2 + c^2 + 2 \left( \frac{1}{a} + \frac{1}{b} + \frac{1}{c} \right) = a^2 + b^2 + c^2 + 2(ab + bc + ca) = (a + b + c)^2.$$

$$\frac{1}{a^2} + \frac{1}{b^2} + \frac{1}{c^2} + 2(a + b + c) = a^2b^2 + b^2c^2 + c^2a^2 + 2abc(a + b + c) = (ab + bc + ca)^2.$$

$$2 \left( \frac{a}{b} + \frac{b}{c} + \frac{c}{a} + \frac{b}{a} + \frac{c}{b} + \frac{a}{c} + 3 \right) = \frac{2(ab(a+b) + bc(b+c) + ca(c+a) + 3abc)}{abc}$$

$$= 2(a + b + c)(ab + bc + ca).$$

By AM-GM Inequality,

$$(a + b + c)^2 + (ab + bc + ca)^2 \geq 2|(a + b + c)(ab + bc + ca)| \geq 2(a + b + c)(ab + bc + ca).$$

This ends the proof. The equality holds for  $a = b = c = 1$ .

▽

**Problem 26 (31, Revised by NguyenDungTN).** *If  $x, y, z$  are positive real numbers, prove that*

$$(x + y + z)^2(yz + zx + xy)^2 \leq 3(y^2 + yz + z^2)(z^2 + zx + x^2)(x^2 + xy + y^2).$$

**Solution 37.** Using the inequality

$$4(a^2 + b^2 + ab) \geq 3(a + b)^2 \quad \forall a, b (\Leftrightarrow (a - b)^2 \geq 0)$$

We have

$$3(y^2 + yz + z^2)(z^2 + zx + x^2)(x^2 + xy + y^2) \geq \frac{4^3}{3^2}(x + y)^2(y + z)^2(z + x)^2.$$

By AM-GM inequality, we get

$$\begin{aligned} 9(x + y)(y + z)(z + x) &= 9(xy(x + y) + yz(y + z) + zx(z + x) + 2xyz) \\ &= 8(xy(x + y) + yz(y + z) + zx(z + x) + 3xyz) + xy(x + y) + yz(y + z) + zx(z + x) - 6xyz \\ &\geq 8(x + y + z)(xy + yz + zx). \end{aligned}$$

So we have the desired result.

▽

**Problem 27 (32, British National Mathematical Olympiad 2007).** *Show that for all positive reals  $a, b, c$*

$$(a^2 + b^2)^2 \geq (a + b + c)(a + b - c)(b + c - a)(c + a - b).$$

**Solution 38 (NguyenDungTN).** Using the familiar inequality

$$xy \leq \frac{(x + y)^2}{4} \quad \forall x, y \in \mathbb{R},$$

we have

$$\begin{aligned} (a + b + c)(a + b - c)(b + c - a)(c + a - b) &= ((a + b)^2 - c^2)(c^2 - (a - b)^2) \\ &\leq \frac{((a + b)^2 - c^2 + c^2 - (a - b)^2)^2}{4} = (a^2 + b^2)^2. \end{aligned}$$

Equality holds when  $(a + b)^2 - c^2 = c^2 - (a - b)^2 \Leftrightarrow c^2 = a^2 + b^2$ .

▽

**Problem 28 (34, Mathlinks, Revised by VanDHHK).** Let  $a, b, c, d$  be real numbers such that  $a^2 \leq 1, a^2+b^2 \leq 5, a^2+b^2+c^2 \leq 14, a^2+b^2+c^2+d^2 \leq 30$  Prove that  $a+b+c+d \leq 10$ .

**Solution 39.** By hypothesis, we have

$$12a^2 + 6b^2 + 4c^2 + 3d^2 \leq 120.$$

By Cauchy-Schwarz Inequality, we have

$$100 = (12a^2 + 6b^2 + 4c^2 + 3d^2) \left( \frac{1}{12} + \frac{1}{6} + \frac{1}{4} + \frac{1}{3} \right) \geq (a + b + c + d)^2$$

Therefore  $a + b + c + d \leq |a + b + c + d| \leq 10$ .

▽